

AD-A126 959

DIELECTRIC DISPERSION IN OPTICAL ELECTRON TRANSFER IN  
SOLUTION(U) NEW YORK UNIV NY DEPT OF CHEMISTRY  
P DELAHAY MAR 83 NYU/DC/TR-2-NEW-SERIES-2

1/


UNCLASSIFIED

N00014-82-K-0113

F/G 20/10

NL

END  
DATE  
FILMED  
DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

6

OFFICE OF NAVAL RESEARCH

Contract N00014-82-K-0113

Task No. NR 359-258

TECHNICAL REPORT NO. NYU/DC/TR-2-NEW-SERIES-2

DIELECTRIC DISPERSION IN OPTICAL ELECTRON TRANSFER IN SOLUTION

by

Paul Delahay

Accepted for publication in

Chemical Physics Letters

New York University  
Department of Chemistry  
New York, NY

March 1983

DTIC  
ELECTE  
APR 19 1983  
S B

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

83 04 19 034

ADA 126959

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DD FORM 1473  
1 JAN 73

Unclassified

**SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)**

# DIELECTRIC DISPERSION IN OPTICAL ELECTRON TRANSFER IN SOLUTION

PAUL DELAHAY

Department of Chemistry, New York University, New York, New York 10003, USA

Received

Dielectric dispersion causes the outer-sphere reorganization free energy to vary with photon energy in optical electron transfer. The resulting distortion of the spectral response for photoelectron emission may yield misleading evidence about the emission yield vs. photon energy relationship. Dispersion is taken into account in the calculation of emission threshold energies and the correlation between optical and thermal electron transfer. Results are given for  $V^{2+}$ ,  $Cr^{2+}$ ,  $Mn^{2+}$ ,  $Fe^{2+}$ ,  $Co^{2+}$ ,  $Fe(CN)_6^{4-}$ .

## 1. INTRODUCTION

Optical electron transfer in solution can be interpreted [1] on the basis of ideas transposed from theories of thermal electron transfer [1-3]. Thus, optical electron transfer is considered to be fast in comparison with subsequent reorganization of nuclear positions. The free energy of reorganization  $R$  is equated to the sum of inner- and outer-sphere contributions,  $R_{in}$  and  $R_{out}$ . These are treated microscopically ( $R_{in}$ , sec. 4) and as non-equilibrium polarization of a continuous medium (sec. 2), respectively. The resulting expression for  $R_{out}$  [1-4] contains the optical dielectric constant  $\epsilon_{op}$  of the solvent. This quantity varies with the photon energy  $E$  at which optical electron transfer occurs because of dielectric dispersion of the solvent. The free energy  $R_{out}$  therefore is a function of  $E$  in contrast to the thermal case in which  $\epsilon_{op}$  is a constant, e.g.,  $\epsilon_{op} = 1.77$  for water at 25°C. The dependence of  $R_{out}$  on  $E$  has significant implications which are examined in the present paper for the case of photoelectron emission by aqueous solutions in the 5 to 11.5 eV range [5].

## 2. FREE ENERGY OF OUTER-SPHERE REORGANIZATION WITH DIELECTRIC DISPERSION

The real and imaginary parts of the dielectric constant of a non-ideally transparent liquid are  $\epsilon_1 = n^2 - k^2$  and  $\epsilon_2 = 2nk$ , respectively. There,  $n$  is the refractive index and  $k$  the absorption coefficient of the medium. The displacement current in such a dielectric is proportional to the real part  $\epsilon_1$ , and consequently the electronic polarization of the medium depends on  $\epsilon_1$  at the appropriate frequency in the optical range. The value of  $n^2 - k^2$  must be taken at the photon energy  $E$  at which electron transfer occurs. Thus, electron transfer is caused by photon absorption, and the electronic polarization of the medium therefore responds to the change of ionic valence of the photoionized species in the same scale of time as photon absorption. The quantity  $n^2 - k^2$  therefore must be substituted for the optical dielectric constant in the expression for  $R_{out}$ . Thus,

$$R_{out}(E) = [(n^2 - k^2)^{-1} - \epsilon_s^{-1}]e^2/2a, \quad (1)$$

where  $\epsilon_s$  is the static dielectric constant of the solvent,  $e$  the electronic charge, and  $a$  the radius of the assumed spherical boundary between inner- and outer-sphere regions. The reduced and oxidized species (denoted respectively by the the subscripts  $r$  and  $o$ ) have somewhat different cavity radii, and one sets  $a = 2a_r a_o / (a_r + a_o)$  as is done for thermal electron transfer [6].

Figure 1 shows the effect of dielectric dispersion on  $R_{out}(E)$  of eq. (1). Results are displayed as the variation of the ratio  $F$  of  $R_{out}(E)$  at  $E$  to the value of  $R_{out}(1.77)$  for  $n^2 = 1.77$  and  $k^2 = 0$  (water at 25°C). Values of  $n$  and  $k$  were taken from [7]. The shape of the curve in fig. 1 is accounted for by the effect of normal and anomalous dispersion (absorption bands at 8.2 and 10.0 eV). It is seen that  $R_{out}(E)$  can be as low as 65 per cent of  $R_{out}(1.77)$  and that  $R_{out}(E)$  changes quite rapidly in rather narrow intervals of  $E$ . Figure 1 applies to pure water, and variations of  $F$  may be somewhat different for concentrated electrolytes, especially in the range of absorption bands of the

anion (charge transfer to the solvent). The variations of  $\epsilon_s$  in (1) with electrolyte concentration are not significant since  $\epsilon_s^{-1}$  for aqueous solutions is small in comparison with  $(n^2 - k^2)^{-1}$ .

### 3. DETERMINATION OF THRESHOLD ENERGIES BY EXTRAPOLATION

Photoelectron emission by solutions is studied [5] by determining the number of electrons collected (on an electrode above the solution) per incident photon as a function of the photon energy  $E$ . The emission yield  $Y$  thus measured is proportional to the emission current density of theory [8] to a good measure under actual experimental conditions. It was shown [8] that  $Y$  is proportional to  $(E - E_t)^p$ , where  $E_t$  is the threshold energy of the species being photoionized and  $p = 2$  or  $5/2$ . This relationship follows from a more general equation for  $Y$  under suitable simplifying conditions. The exponent  $p = 2$  corresponds to the normal case in which image forces are taken into account at the solution-vacuum interface. The value  $p = 5/2$  holds if one assumes that the image forces are negligible.

The threshold energy  $E_t$  is determined from a plot of  $Y^{1/p}$  against  $E$ . This procedure was applied to water [9], inorganic anions [10] and cations [11], and weak acids and bases [12] in aqueous solution. It was found that the exponent  $p = 5/2$  gave better fits to linear plots than  $p = 2$  for species with  $E_t < 8$  eV. Conversely, the exponent  $p = 2$  was preferable to  $5/2$  for  $E_t \geq 8$  eV. The statistical evidence for these values of  $p$  was fairly unambiguous in most cases, but the underlying reasons for the choice of  $p = 5/2$  were mystifying. The absence of image forces is difficult to account for, and anyhow the magnitude of image forces would conceivably depend on the kinetic energy of mobile electrons in the liquid and not on the absolute value of the photon energy. These puzzling results are accounted for by the effect of dispersion on  $R_{out}(E)$  as will now be shown.



Availability Codes	
Dist	Avail and/or Special
<b>A</b>	

The threshold energy is the sum of several contributions one of which is the free energy of outer-sphere reorganization  $R_{\text{out}}$  (sec. 4). Call  $E_t^0$  the threshold energy at  $Y = 0$  in the linear extrapolation plot of  $Y^{1/p}$  against  $E$ . One has  $R_{\text{out}} = R_{\text{out}}^0$  at  $E = E_t^0$ , and a different  $R_{\text{out}}(E)$  at  $E > E_t^0$ . The value of  $E_t$  in  $(E - E_t)^p$  therefore is not the constant  $E_t^0$  but the quantity,

$$E_t = E_t^0 - [R_{\text{out}}^0 - R_{\text{out}}(E)], \quad (2)$$

depending on  $E$ . Since  $R_{\text{out}}(E)$  may differ from  $R_{\text{out}}^0$  by several tenths of electronvolt, plots of  $Y^{1/p}$  against  $E$  are distorted.

Variations of  $(E - E_t)^2$  with  $E - E_t^0$  are shown in Fig. 2 for  $E_t^0 = 6$  and 8 eV, respectively. Values of  $E - E_t$  were calculated from fig. 1 and the values of  $R_{\text{out}}^0 = 1.02$  and 0.81 eV at  $E_t^0 = 6$  and 8 eV, respectively. These values of  $R_{\text{out}}^0$  correspond to the typical case of  $R_{\text{out}}(1.77) = 1.15$  eV for the cations of sec. 4. The curve  $(E - E_t)^2$  vs.  $(E - E_t^0)$  for  $E_t^0 = 6$  eV in Fig. 2 is quite close to the points representing  $(E - E_t^0)^{5/2}$ . This result follows directly from Fig. 1. The ratio  $F$  decreases quite rapidly with increasing  $E$  in the 6 to 7.6 eV range, and consequently  $(E - E_t)^2$  increases more rapidly than  $(E - E_t^0)^2$ . Conversely, the curve  $(E - E_t)^2$  vs.  $E - E_t^0$  for  $E_t^0 = 8$  eV in Fig. 2 is fairly close to the quadratic dependence  $(E - E_t^0)^2$  because the decrease of  $F$  between  $E = 8.6$  and 9.4 eV is less pronounced than the variations of  $F$  for  $E_t^0 = 6$  eV. Apparent variation of  $p$  from  $5/2$  to 2 for the results of Fig. 2 are solely due to the change of  $R_{\text{out}}(E)$  with  $E$  as a result of dispersion. The statistical evidence in [10-12] upon which the choice of  $p = 5/2$  was based therefore is misleading and does not provide any proof that the  $(E - E_t)^{5/2}$  relationship is in fact observed. A more elaborate analysis must be developed on the basis of the quadratic dependence regardless of photon energy and with consideration of the variations of  $R_{\text{out}}(E)$  with photon energy.



#### 4. CALCULATED FREE ENERGIES OF EMISSION VERSUS EXPERIMENTAL THRESHOLD ENERGIES

The free energy  $\Delta G_m$  for photoelectron emission by an aqueous solution of a donor  $D(\text{aq})$  is [5,11]



$$\Delta G_m = \Delta G_H + \Delta G + R + |e|\Delta x. \quad (3)$$

There,  $\Delta G_H$  and  $\Delta G$  are the changes of free energy for the reactions,



respectively;  $R$  is the reorganization free energy taken as a positive quantity for convenience; and  $\Delta x$  is the difference between the surface potentials  $\chi[D(aq)]$  of the solution of  $D(aq)$  and  $\chi(H_2O)$  of water.

One has [13]  $\Delta G_H = 4.48 \pm 0.05$  eV on the basis of a current determination [14] of the real free energy of hydration ( $-11.25$  eV) of the proton. This value of  $\Delta G_H$  includes the contribution of the surface potential  $\chi(H_2O)$  of water. The latter cannot be measured but can be estimated ( $0.08 \pm 0.06$  eV [15]). The surface potential  $\chi[D(aq)]$  is generally different from that of water, and the experimentally measurable [15] difference  $\Delta x$  therefore appears in (3). Since the surface potential pertains to the change of potential from the liquid to vacuum, a positive value of  $\Delta x$  corresponds to a higher barrier to emission than for  $\Delta x = 0$ . Hence, the quantity  $|e|\Delta x$  in (3) is preceded by a positive sign. The correction is quite minor anyhow since  $|e|\Delta x$  is in general smaller than 0.05 to 0.1 eV in absolute value [15].

The reorganization free energy is  $R = R_{out}(E) + R_{in}$ , where  $R_{in}$  is computed from the bond-stretching model used for thermal electron transfer [1]. Thus,

$$R_{in} = (N/2)f(\Delta q_0)^2, \quad (4)$$

where  $N$  is the number of ligands of the donor  $D(aq)$ ;  $\Delta q_0$  is the change of equilibrium positions for vibration from the reduced species  $D(aq)$  to  $D^+(aq)$ ; and  $f$  is the reduced force constant [6],

$$f = 2f_r f_o / (f_r + f_o). \quad (5)$$

Equation (4) gives an energy and not a free energy [1,6], but the resulting error can be neglected since the model upon which eq. (4) is based is somewhat simplified for hexaquo ions, for instance.

The  $f$ -values from [16] previously used [4] were computed from a Urey-Bradley potential in [17]. This gives  $f$ -values which are definitely too low, as was pointed out [18] to the writer, and experimental force constants must be used. One has [6],

$$f_r = 4\pi^2 \nu_r^2 c^2 \mu, \quad (6)$$

where  $\nu_r$  is the stretching frequency (in  $\text{cm}^{-1}$ ) and  $\mu$  the reduced mass. A similar equation applies to  $f_0$ . Two limiting cases have been considered for  $\mu$  in the case of hexaquo ions discussed below: (i)  $\mu$  is set equal to the mass of a single water molecule  $m_{\text{H}_2\text{O}}$  [6,19]; (ii)  $\mu = m_{\text{H}_2\text{O}} m_M / (m_{\text{H}_2\text{O}} + m_M)$ , where  $m_M$  is the atomic mass of  $M^{2+/3+}$  [20]. Method (i) slightly overestimates  $\mu$ . Method (ii) certainly underestimates  $\mu$  (by  $\approx 20$  to 25 per cent for the cations considered here) since it presupposes vibration between a "dry" cation  $M^{2+/3+}$  and a water molecule. The first method was applied here. In the case of  $\text{Fe}(\text{CN})_6^{4-/3-}$ ,  $\mu$  was set equal to the CN mass.

The  $\Delta q_0$ 's in (4) computed from crystallographic metal-ligand distances from X-ray diffraction [6,19,20] generally agree with the  $\Delta q_0$ 's recently determined by EXAFS for solutions [19]. The  $\Delta q_0$  for  $\text{Fe}(\text{CN})_6^{4-/3-}$  is particularly low: 0.026 Å (crystallographic) vs. 0.01 Å (EXAFS, solutions) [19]. The calculation of  $R_{in}$  is sensitive to  $\Delta q_0$ , and an error of  $\pm 0.01$  Å on this quantity affects  $R_{in}$  by  $\approx \pm 0.1$  to  $\pm 0.15$  eV for the cations discussed below.

Values of  $\Delta G_m$  computed from eq. (3) are listed in Table 1 with the corresponding  $E_t$ 's. Agreement can be considered to be good except for  $\text{Fe}^{2+}$  for an a priori (no data from emission) calculation of  $\Delta G_m$  in view of the uncertainties on calculated  $R_{in}$ 's and extrapolated  $E_t$ 's. The agreement for  $\text{Fe}(\text{CN})_6^{4-}$  shows that the threshold energy (5.7 eV) can be assigned to a direct bound-continuum transition. The threshold was attributed to autoionization of a bound state in [23] on the basis of the low value of  $E_t = 5.5$  eV (obtained by extrapolation with  $p = 5/2$ ) and a higher  $R_{in}$  value from

[16]. Assignment of emission autoionization bands [23], in general, will have to be reexamined in the light of the present work. The discrepancy for  $\text{Fe}^{2+}$  is ascribed mostly to the high  $E_t$ 's (for  $p = 5/2$  and  $p = 2$ ) in Table 1 rather than the calculated  $R_{in}$ . The latter indeed agrees well with the value  $R_{in} = 0.69$  eV computed from experimental data on thermal electron transfer in sec. 5. Such a calculation from kinetic data is reliable for  $\text{Fe}^{2+/3+}$  [6].

## 5. CORRELATION BETWEEN OPTICAL AND THERMAL ELECTRON TRANSFER

Optical and thermal electron transfer were correlated in [4] without consideration of dielectric dispersion. It was shown that the outer-sphere reorganization free energy  $R_{out}^X$  for electron exchange (e.g., between  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$ ) is practically the same as  $R_{out}$  for photoelectron emission. The inner-sphere term  $R_{in}^X$  for exchange is twice  $R_{in}$  for emission since two ions are involved in exchange and only one in emission. Thus,  $R_{out}^X = R_{out}$  and  $R_{in}^X = 2R_{in}$ , and the reorganization free energy for exchange is  $R^X = R + R_{in}$ . The effect of dielectric dispersion will now be considered.

The quantity  $R_{out}^X$  for exchange is generally computed for  $\epsilon_{op} = 1.77$  for dilute aqueous solutions at  $25^\circ\text{C}$ , that is,  $R_{out}^X = R_{out}(1.77)$ . Hence,

$$R^X = R + R_{in} + \Delta R_{out}, \quad (7)$$

where the correction for dispersion is  $\Delta R_{out} = R_{out}(1.77) - R_{out}(E)$ ,  $R_{out}(E)$  being computed at  $E = E_t$ . The free energy of activation  $\Delta G^\ddagger$  for electron exchange without net change of free energy is [1-3],  $\Delta G^\ddagger = w + R^X/4$ , where  $w$  is the work required to bring from infinity in solution the two reactants together in the activated complex.

Values of  $\Delta G^\ddagger$  from (7) are listed in Table 2 with experimental values taken from [1]. This comparison, it must be stressed, is only tentative because calculated values of  $\Delta G^\ddagger$  should be compared with the activation free energy for electron transfer within the precursor complex [6]. Data are included only for couples for which  $\Delta G^\ddagger$  can be computed reasonably well from  $\Delta G^\ddagger = R^X/4 + w$  [1-3]. Thus, exchange for  $\text{Co}^{2+/3+}$  is known to be anomalously slow [19]. The

"experimental" value  $\Delta G^\ddagger = 0.75$  eV for  $\text{Mn}^{2+/3+}$  obtained in [1] from the Marcus cross relationship differs markedly from the value  $\Delta G^\ddagger = 1.06$  eV computed from  $R^X$ . Experimental (0.47 eV [1]) and calculated (0.31 eV)  $\Delta G^\ddagger$ 's for  $\text{Fe}(\text{CN})_6^{4-/3-}$  also disagree.

Discrepancies in Table 2 can be interpreted on the basis of values of  $R_{\text{in}}$  computed from  $\Delta G^\ddagger = R^X/4 + w$  [1-3] by using the experimental  $\Delta G^\ddagger$ 's in Table 2 and calculated values of  $R_{\text{out}}^X (= R_{\text{out}}(1.77))$ . Errors on  $\Delta G^\ddagger$  and  $R_{\text{out}}^X$  affect the value of  $R_{\text{in}}$ , and application of the equation for  $\Delta G^\ddagger$  is assumed to be justified. Values of  $R_{\text{in}}$  thus obtained obviously must be viewed with caution. One has  $R_{\text{in}} = 1.09, 1.39, 0.69$  eV for  $\text{V}^{2+}, \text{Cr}^{2+}, \text{Fe}^{2+}$ , respectively, vs. 1.41, 1.27, 0.73 eV in Table 1. One concludes that the main source of discrepancy between calculated and experimental  $\Delta G^\ddagger$ 's in Table 2 is mostly the high  $R_{\text{in}}$  of Table 1 for  $\text{V}^{2+/3+}$  and the high  $E_t$  (and consequently high  $R$ ) for  $\text{Fe}^{2+/3+}$ . Agreement for  $\text{Cr}^{2+/3+}$  is reasonable.

In conclusion, consideration of dispersion is essential in future determinations of experimental threshold energies. Free energies of emission computed from data independent of emission generally agree with available threshold energies within extrapolation error. Correlation between optical and thermal electron transfer provides, whenever feasible, further verification of the self-consistency of all data.

#### ACKNOWLEDGMENT

This work was supported by the Office of Naval Research and the National Science Foundation. The author thanks his colleagues R. Parsons (Bellevue, France) and N. Sutin (Brookhaven, USA) for the communication of manuscripts prior to publication.

#### REFERENCES

- [1] R. D. Cannon, Electron transfer reactions (Butterworths, London, 1980) pp. 175-222, 267-297.

- [2] R. A. Marcus, J. Chem. Phys. 24 (1956) 966, 979; 43 (1965) 679.
- [3] N. S. Hush, Trans. Faraday Soc. 57 (1961) 557.
- [4] P. Delahay, Chem. Phys. Letters 87 (1982) 607.
- [5] P. Delahay, Accounts Chem. Res. 15 (1982) 40.
- [6] B. S. Brunshwig, J. Logan, M. D. Newton, N. Sutin, J. Am. Chem. Soc. 102 (1980) 5798.
- [7] J. M. Heller, Jr., R. N. Hamm, R. D. Birkhoff, L. R. Painter, J. Chem. Phys. 60 (1974) 3483.
- [8] A. M. Brodsky and A. V. Tsarevsky, J. Chem. Soc., Faraday Trans. II 72 (1976) 1781.
- [9] P. Delahay and K. von Burg, Chem. Phys. Letters 83 (1981) 250.
- [10] K. von Burg and P. Delahay, Chem. Phys. Letters 78 (1981) 287.
- [11] P. Delahay, K. von Burg, A. Dziedzic, Chem. Phys. Letters 79 (1981) 157.
- [12] K. von Burg and P. Delahay, Chem. Phys. Letters 86 (1982) 528.
- [13] R. Parsons, private communication.
- [14] J. R. Farrell and P. McTigue, J. Electroanal. Chem. 139 (1982) 37.
- [15] J. E. B. Randles, Phys. Chem. Liq. 7 (1977) 107.
- [16] J. M. Hale, in Reactions of molecules at electrodes, ed. N. S. Hush (Wiley-Interscience, New York, 1971) pp. 229-257.
- [17] I. Nakagawa and T. Shimanouchi, Spectrochim. Acta 20 (1964) 429.
- [18] J. O'M. Bockris, private communication.
- [19] B. S. Brunshwig, C. Creutz, D. H. Macartney, T-K. Sham, N. Sutin, Disc. Faraday Soc., in press.
- [20] J. O'M. Bockris, S. U. M. Khan, D. B. Matthews, J. Res. Inst. Catalysis, Hokkaido 22 (1974) 1.
- [21] N. Sutin, Ann. Rev. Nucl. Sci. 12 (1962) 285.
- [22] K. Nakamoto, Infrared and Raman spectra of inorganic and coordination compounds, 3rd ed. (Wiley-Interscience, New York, 1978), pp. 228, 264.
- [23] P. Delahay, Chem. Phys. Letters 90 (1982) 425.

Table 1

Calculated free energies of emission vs. experimental threshold energies.

	$\Delta G$ (eV)	$R_{out}$ <sup>a)</sup> (eV)	$R_{in}$ <sup>b)</sup> (eV)	$\Delta G_m$ (eV)	$E_t$ <sup>c)</sup> (eV)
$V^{2+}$	-0.25	0.99	1.41	6.63	6.38 (p=5/2) 6.74 (p=2)
$Cr^{2+}$	-0.41	1.01	1.27	6.35	6.14 (p=5/2) 6.51 (p=2)
$Mn^{2+}$	1.56	0.95	1.51	8.50	8.08 (p=2)
$Fe^{2+}$	0.77	0.93	0.73	6.91	7.35 (p=5/2) 7.47 (p=2)
$Co^{2+}$	1.84	0.95	1.63	8.90	8.60 (p=2)
$Fe(CN)_6^{4-}$	0.36	0.77	0.06	5.67	5.5 (p=5/2) 5.7 (p=2)

a) From  $FR_{out}(1.77)$  with  $F$  at  $E = \Delta G_m$  (Fig. 1).

Values of

$R_{out}(1.77)$  computed, respectively, for  $a = 3.48, 3.48, 3.56, 3.51, 3.48$  Å from  $a_r$  and  $a_o$  in [21] and  $a = 4.65$  Å from [16].

b) Calculated for  $N = 6$ ;  $f = 1.7 \times 10^5$  dyne  $cm^{-1}$  for  $Cr^{2+}$  [19];  $f$  from  $\nu_r = 389$   $cm^{-1}$  and  $\nu_o = 490$   $cm^{-1}$  for all other cations except  $\nu_r = 395$   $cm^{-1}$  for  $Mn^{2+}$  [20,22];  $f = 4.5 \times 10^5$  dyne  $cm^{-1}$  for  $Fe(CN)_6^{4-}$  from  $\nu_r = 585$   $cm^{-1}$  and  $\nu_o = 511$   $cm^{-1}$  from [20,22];  $\Delta q_0 = 0.20$  ( $Cr^{2+}$ ),  $0.14$  ( $Fe^{2+}$ ),  $0.21$  ( $Co^{2+}$ ),  $0.026$  ( $Fe(CN)_6^{4-}$ ) Å from [19] and  $0.195$  ( $V^{2+}$ ),  $0.2$  ( $Mn^{2+}$ ) Å from [20].

c) First  $E_t$  for  $p = 5/2$  for  $V^{2+}$ ,  $Cr^{2+}$ ,  $Fe^{2+}$ ,  $Fe(CN)_6^{4-}$ . Second  $E_t$  from emission spectra used in [11] for all cations and [23] for  $Fe(CN)_6^{4-}$ , but for extrapolation with  $p = 2$ . Only extrapolation for  $p = 2$  for  $Mn^{2+}$  and  $Co^{2+}$  [11].

Table 2

Calculated and experimental free energies of activation for electron exchange

	$R^a)$	$\Delta R_{out}^b)$	$w^c)$	$\Delta G^\ddagger^d)$	$\Delta G^\ddagger^e)$
	(eV)	(eV)	(eV)	(calculated) (eV)	(experimental) (eV)
$V^{2+/3+}$	2.15 (p=5/2)	0.15	0.04	0.97 (p=5/2)	0.87
	2.51 (p=2)			1.06 (p=2)	
$Cr^{2+/3+}$	2.07 (p=5/2)	0.13	0.05	0.92 (p=5/2)	1.03
	2.44 (p=2)			1.01 (p=2)	
$Fe^{2+/3+}$	2.10 (p=5/2)	0.18	0.06	0.81 (p=5/2)	0.69
	2.22 (p=2)			0.84 (p=2)	

a) From eq. (3) and data in Table 1.

b) From  $R_{out}$  in Table 1 and the corresponding  $R_{out}(1.77)$ .

c) From [4].

d) For  $R_{in}$  in Table 1.

e) From [1].

## Captions to Figures

Fig. 1. Plot of the ratio  $F$  of the outer-sphere reorganization free energy  $R_{\text{out}}(E)$  at the photon energy  $E$  to  $R_{\text{out}}(1.77)$  computed for  $n^2 = 1.77$  and  $k^2 = 0$ .

Fig. 2. Spectral response for emission  $(E - E_t)^2$  with consideration of dispersion compared with the responses  $(E - E_t^0)^2$  and  $(E - E_t^0)^{5/2}$  without dispersion.






FIG. 1




FIG. 2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	<u>No. Copies</u>		<u>No. Copies</u>
Professor K. Wilson Department of Chemistry, B-014 University of California, San Diego La Jolla, California 92093	1	Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1
Professor C. A. Angell Department of Chemistry Perdue University West Lafayette, Indiana 47907	1	Dr. B. Vonnegut State University of New York Earth Science Building 1400 Washington Avenue Albany, New York 12203	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SOO 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Chemistry & Materials Science Department Laurence Livermore National Lab. Livermore, California 94550	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auburn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research and Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Johnson Controls 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles and Space Co., Inc. P. O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attn: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1		
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. H. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139		Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
S. Ruby DOE (STOR) 600 E Street Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton 509 5NH England	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. Bernard Spielvogel U. S. Army Research Office P. O. Box 12211 Research Triangle Park, NC 27709	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	Dr. Denton Elliott Air Force Office of Scientific Research Bolling AFB Washington, D.C. 20332	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. David Aikens Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. A. P. B. Lever Chemistry Department York University Downsview, Ontario M3J1P3 Canada	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England		Dr. Stanislaw Szpak Naval Ocean Systems Center Code 6343 San Diego, California 95152	1
Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362		Dr. Gregory Farrington Department of Materials Science and Engineering University of Pennsylvania Philadelphia, Pennsylvania 19104	
Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, D.C. 20376		Dr. Bruce Dunn Department of Engineering & Applied Science University of California Los Angeles, California 90024	

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
M. L. Robertson Manager, Electrochemical and Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1	Dr. D. Cipris Allied Corporation P. O. Box 3000R Morristown, New Jersey 07960	1
Dr. Micha Tomkiewicz Department of Physics Brooklyn College Brooklyn, New York 11210	1	Dr. M. Philpot IBM Corporation 5600 Cottle Road San Jose, California 95193	1
Dr. Lesser Blum Department of Physics University of Puerto Rico Rio Piedras, Puerto Rico 00931	1	Dr. Donald Sandstrom Washington State University Department of Physics Pullman, Washington 99164	1
Dr. Joseph Gordon, II IBM Corporation K33/281 5600 Cottle Road San Jose, California 95193	1	Dr. Carl Kannewurf Northwestern University Department of Electrical Engineering and Computer Science Evanston, Illinois 60201	1
Dr. Robert Somoano Jet Propulsion Laboratory California Institute of Technology Pasadena, California 91103	1	Dr. Edward Fletcher University of Minnesota Department of Mechanical Engineering Minneapolis, Minnesota 55455	1
Dr. Johann A. Joebstl USA Mobility Equipment R&D Command DRDME-EC Fort Belvoir, Virginia 22060	1	Dr. John Fontanella U.S. Naval Academy Department of Physics Annapolis, Maryland 21402	1
Dr. Judith H. Ambrus NASA Headquarters M.S. RTS-6 Washington, D.C. 20546	1	Dr. Martha Greenblatt Rutgers University Department of Chemistry New Brunswick, New Jersey 08903	1
Dr. Albert R. Landgrebe U.S. Department of Energy M.S. 6B025 Forrestal Building Washington, D.C. 20595	1	Dr. John Wassib Kings Mountain Specialties P. O. Box 1173 Kings Mountain, North Carolina 28086	1



TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>	<u>No. Copies</u>
Dr. J. J. Brophy University of Utah Department of Physics Salt Lake City, Utah 84112	1	
Dr. Walter Roth Department of Physics State University of New York Albany, New York 12222	1	
Dr. Thomas Davis National Bureau of Standards Polymer Science and Standards Division Washington, D.C. 20234	1	
Dr. Charles Martin Department of Chemistry Texas A&M University	1	
Dr. Anthony Sammells Institute of Gas Technology 3424 South State Street Chicago, Illinois 60616	1	
Dr. H. Tachikawa Department of Chemistry Jackson State University Jackson, Mississippi 39217	1	
Dr. W. M. Risen Department of Chemistry Brown University Providence, Rhode Island	1	

**DAT**  
**ILM**